arXiv:1601.01567v1 [math.DG] 7 Jan 2016

The Intersection of a Hyperplane with a Lightcone in the Minkowski Spacetime

Pengyu Le

Abstract

Klainerman, Luk and Rodnianski derived an anisotropic criterion for formation of trapped surfaces in vacuum, extending the original trapped surface formation theorem of Christodoulou. The effort to understand their result led us to study the intersection of a hyperplane with a lightcone in the Minkowski spacetime. For the intrinsic geometry of the intersection, depending on the hyperplane being spacelike, null or timelike, it has the constant positive, zero or negative Gaussian curvature. For the extrinsic geometry of the intersection, we find that it is a noncompact marginal trapped surface when the hyperplane is null. In this case, we find a geometric interpretation of the Green's function of the Laplacian on the standard sphere. In the end, we contribute a clearer understanding of Klainerman, Luk and Rodnianski's result.

1 Introduction

In 1965 work of Penrose [3], he introduced the concept of a closed trapped surface, i.e. a closed spacelike surface with the property that the area element decreases pointwise for any infinitesimal displacement along the future null direction normal to the surface. Based on this concept, Penrose proved his famous incompleteness theorem. The incompleteness theorem says the following: a vacuum spacetime cannot be future null geodesically complete if it contains a non-compact Cauchy hypersurface and moreover contains a closed trapped surface. Then it follows that the spacetime with a complete null infinity which contains a closed trapped surface must contain a future event horizon the interior of which contains the closed trapped surface.

The incompleteness theorem gives rise to the following problem: could the closed trapped surface be formed dynamically in vacuum? In the important monograph of Christodoulou [1], he answered this problem positively by showing that the closed trapped surface could be formed by sufficiently strong incoming gravitational waves. Christodoulou introduced a new method called "short pulse method", which allows him to establish an existence theorem for a development of initial data which is large enough so that the trapped surfaces have a chance to form within this development. In the proof, he adopted a two parameter family of surfaces, which are the intersections of the incoming family with the outgoing family of null hypersurfaces. Then provided energy of the incoming gravitation waves being large, the intersection of the last incoming null hypersurface with the later outgoing null hypersurface is trapped.

Klainerman, Luk and Rodnianski derived an anisotropic criterion for formation of trapped surfaces in vacuum in [2] , using the existence theorem of Christodoulou. The original trapped surface formation theorem requires a lower bound on energy per solid angle of the incoming gravitational waves in all directions. They showed instead that, if the energy per solid angle of the incoming gravitational waves in some neighbourhood of some direction is sufficiently large depending on the size of the neighbourhood, then the development of initial data contains a trapped surface. Although none in the two parameter family of surfaces adopted in the existence theorem is trapped, they could find another closed spacelike surface which is trapped in the last incoming null hypersurface.

The result of Klainerman, Luk and Rodnianski is surprising since the spacetime could be flat in a large part of the development of initial data they considered. We find that the intersection of a null hyperplane with a lightcone in the Minkowski spacetime is actually marginal trapped. As an application, we give a easier way to find a trapped surface.

2 The geometry of a spacelike surface in a vacuum spacetime

We consider a spacelike surface S in a vacuum spacetime (M,g). On the normal bundle $\mathrm{N}S$ of S, we can introduce a null frame $\{L,\underline{L}\}$. $L,\underline{L}$ are future directed null vector fields in $\mathrm{N}S$ and satisfies

$$g(L,\underline{L}) = -2. \tag{2.1}$$

Such choice of a null frame is not unique, since for any positive function a on S, $\{aL, a^{-1}\underline{L}\}$ is another null frame. Associated with a null frame $\{L,\underline{L}\}$, we can define the following 2-covariant tensor fields $\chi,\underline{\chi}$ and the1-form ζ over S by

$$\chi(X,Y) = g(\nabla_X L, Y), \quad \underline{\chi}(X,Y) = g(\nabla_X \underline{L}, Y), \quad \zeta(X) = \frac{1}{2} g(\nabla_X L, \underline{L}) \tag{2.2}$$

for any vector fields X,Y tangential to S.

χ is the second fundamental form of S corresponding to L and $\underline{\chi}$ is the second fundamental form corresponding to $\underline{L}$. We call ζ the torsion of the null frame $\{L,\underline{L}\}$. The traces $\mathrm{tr}\chi, \mathrm{tr}\underline{\chi}$ of $\chi,\underline{\chi}$ are called the null expansions and the tracefree parts $\widehat{\chi},\widehat{\underline{\chi}}$ are called the shears.

Definition 2.1.
1. A spacelike surface S is called trapped if the two null expansions are both negative, i.e. $\mathrm{tr}\chi < 0$ and $\mathrm{tr}\underline{\chi} < 0$.
2. A spacelike surface S is called marginally trapped if one of the null expansion is identically zero and the other is negative.

We choose an orthogonal frame $\{e_1,e_2\}$ on the tangent bundle $\mathrm{T}S$ such that $\{e_1,e_2,e_3 = \underline{L}, e_4 = L\}$ is a positive oriented frame on (M,g). We denote the area element of S by $\not{\epsilon}$. Let the

Latin index $a, b, \cdots$ denote $1, 2$. We define the 2-covariant S-tensor fields $\alpha, \underline{\alpha}$, the S-tangential 1-forms $\beta, \underline{\beta}$ and the functions ρ, σ on S as following:

$$\alpha_{ab} = \mathrm{R}_{4ab4}, \quad \underline{\alpha}_{ab} = \mathrm{R}_{3ab3}, \tag{2.3}$$

$$\beta_a = \tfrac{1}{2}\mathrm{R}_{a443}, \quad \underline{\beta}_a = \frac{1}{2}\mathrm{R}_{a343}, \tag{2.4}$$

$$\rho = \tfrac{1}{4}\mathrm{R}_{4334}, \quad \sigma \cdot \not{\epsilon}_{ab} = \frac{1}{2}\mathrm{R}_{ab43}. \tag{2.5}$$

Because of the symmetries of the curvature tensor and the Ricci flat condition, we have

$$\mathrm{R}_{abc3} = -\not{\epsilon}_{ab}\not{\epsilon}_c{}^d\underline{\beta}_d, \quad \mathrm{R}_{abc4} = \not{\epsilon}_{ab}\not{\epsilon}_c{}^d\beta_d \tag{2.6}$$

$$\mathrm{R}_{abcd} = \rho \cdot \not{\epsilon}_{ab}\not{\epsilon}_{cd}, \quad \mathrm{R}_{3ab4} = -\rho \cdot \not{g}_{ab} + \sigma \cdot \not{\epsilon}_{ab}. \tag{2.7}$$

Hence $\{\alpha, \underline{\alpha}, \beta, \underline{\beta}, \rho, \sigma\}$ contains all the curvature components.

We have the following Gauss equation:

$$K + \frac{1}{4}\mathrm{tr}\chi\,\mathrm{tr}\underline{\chi} - \frac{1}{2}(\widehat{\chi}, \widehat{\underline{\chi}}) = -\rho. \tag{2.8}$$

Here

$$(\widehat{\chi}, \widehat{\underline{\chi}}) = (\not{g}^{-1})^{ac}\,(\not{g}^{-1})^{bd}\,\widehat{\chi}_{ab}\,\widehat{\underline{\chi}}_{cd}. \tag{2.9}$$

3 The sections of a lightcone in the Minkowski spacetime $(\mathbb{M}, \eta)$

The Minkowski spacetime is the 4-dim affine space $\mathbb{M}$ endowed with the flat Lorentzian metric η. Let $\{x_0, x_1, x_2, x_3\}$ be the rectangular coordinate system on $\mathbb{M}$ such that $\eta = -\mathrm{d}x_0^2 + \mathrm{d}x_1^2 + \mathrm{d}x_2^2 + \mathrm{d}x_3^2$. The lightcone issuing from the origin o is the subset $C_o = \{-x_0^2 + x_1^2 + x_2^2 + x_3^2 = 0\}$.

Let $\{t, r, \vartheta, \phi\}$ be the spatial polar coordinate system $\{t, r, \vartheta, \phi\}$ where $t \in \mathbb{R}$, $r \in \mathbb{R}_{>0}$, $\vartheta \in (0, \pi)$ and $\phi \in (0, 2\pi)$. The coordinate transformation from the spatial polar coordinates to the rectangular coordinates is

$$(x_0, x_1, x_2, x_3) = (t, r\sin\vartheta\cos\phi, r\sin\vartheta\sin\phi, r\cos\vartheta). \tag{3.1}$$

We define two optical functions $u = \frac{1}{2}(t - r)$ and $v = \frac{1}{2}(t + r)$. Then denote $\{C_u\}$ the level sets of u and $\{\underline{C}_v\}$ the level sets of v. C_u is the future half lightcone issuing from the point $(u, 0, 0, 0)$ and $\underline{C}_v$ is the past half lightcone issuing from the point $(v, 0, 0, 0)$. We denote $S_{u,v}$ the intersection C_u with $\underline{C}_v$.

$\{u, v, \vartheta, \phi\}$ would also be a coordinate system on $\mathbb{M}$ and we call it a double null coordinate system. The metric η takes the form $\eta = -2(\mathrm{d}u \otimes \mathrm{d}v + \mathrm{d}v \otimes \mathrm{d}u) + r^2(\mathrm{d}\vartheta^2 + \sin^2\vartheta\mathrm{d}\phi^2)$ with $r = v - u$. The intrinsic geometry of $S_{u,v}$ is a round sphere of radius $v - u$.

The coordinate vector fields ∂_v and ∂_u are two geodesic null vector fields generating the lightcones C_u and $\underline{C}_v$. Moreover they are normal to the surface $S_{u,v}$ and $\eta(\partial_u, \partial_v) = -2$. Hence $\{\partial_u, \partial_v\}$

is a null frame on $S_{u,v}$. Let us denote $\{\partial_v, \partial_u\}$ by $\{L, \underline{L}\}$. Then we can apply the constructions in the section 2. We have

$$\mathrm{tr}\chi = \frac{2}{r} = \frac{2}{v-u}, \quad \mathrm{tr}\underline{\chi} = -\frac{2}{r} = -\frac{2}{v-u}, \quad \hat{\chi} = 0, \quad \hat{\underline{\chi}} = 0, \quad \zeta = 0, \tag{3.2}$$

and

$$\alpha = 0, \quad \underline{\alpha} = 0, \quad \beta = 0, \quad \underline{\beta} = 0, \quad \rho = 0, \quad \sigma = 0. \tag{3.3}$$

We consider the lightcone $\underline{C}_0$ which is the past half lightcone issuing from the origin. $\{u, \vartheta, \phi\}$ is a coordinate system on $\underline{C}_0$. The metric η restricted to $\underline{C}_0$ takes the form $\eta|_{\underline{C}_0} = u^2(\mathrm{d}\vartheta^2 + \sin^2\vartheta \mathrm{d}\phi^2)$.

Let S be a spacelike surface in $\underline{C}_0$, we could represent it as a graph of a positive function f on the sphere: $S = \{(u = -f(\vartheta,\phi), \vartheta, \phi)\}$. Conversely, for any positive function f on the sphere, we could construct a spacelike surface S_f in $\underline{C}_0$ by $S_f = \{(u = -f(\vartheta,\phi), \vartheta, \phi)\}$. The metric η restricted to S_f takes the form $\eta|_{S_f} = f^2(\vartheta,\phi)(\mathrm{d}\vartheta^2 + \sin^2\vartheta \mathrm{d}\phi^2)$, which is a conformal deformation of the round sphere.

We investigate the geometry of the surface S_f constructed above. The tangent space of S_f is spanned by $\{\widetilde{\partial}_\vartheta, \widetilde{\partial}_\phi\}$ where

$$\widetilde{\partial}_\vartheta = \partial_\vartheta - f_\vartheta \partial_u, \quad \widetilde{\partial}_\phi = \partial_\phi - f_\phi \partial_v. \tag{3.4}$$

$\underline{L} = \partial_u$ is a null vector field normal to S_f. By solving the equation

$$\eta(\widetilde{L}, \widetilde{\partial}_\vartheta) = \eta(\widetilde{L}, \widetilde{\partial}_\phi) = 0, \quad \eta(\widetilde{L}, \underline{L}) = -2, \tag{3.5}$$

we get $\widetilde{L} = \partial_v - 2\eta^{\vartheta\vartheta} f_\vartheta \partial_\vartheta - 2\eta^{\phi\phi} f_\phi \partial_\phi$. Hence $\{\widetilde{L}, \underline{\widetilde{L}}\}$ is a null frame on the normal bundle of S_f where

$$\underline{\widetilde{L}} = \underline{L} = \partial_u, \quad \widetilde{L} = \partial_v - 2\eta^{\vartheta\vartheta} f_\vartheta \partial_\vartheta - 2\eta^{\phi\phi} f_\phi \partial_\phi. \tag{3.6}$$

We can calculate the null expansions $\mathrm{tr}\widetilde{\chi}$ and $\mathrm{tr}\underline{\widetilde{\chi}}$, the shears $\widehat{\widetilde{\chi}}$ and $\widehat{\underline{\widetilde{\chi}}}$, the torsion $\widetilde{\zeta}$ and the curvature components $\widetilde{\alpha}, \underline{\widetilde{\alpha}}, \widetilde{\beta}, \underline{\widetilde{\beta}}, \widetilde{\rho}, \widetilde{\sigma}$ along S_f corresponding to $\{\widetilde{L}, \underline{\widetilde{L}}\}$. Since the Minkowski spacetime is flat, the curvature components are all zero. For our purpose, we are more concerning with the null expansions,

$$\mathrm{tr}\underline{\widetilde{\chi}} = -\frac{2}{f}, \tag{3.7}$$

$$\begin{aligned} \mathrm{tr}\widetilde{\chi} &= \frac{2}{f} - 2\Delta_{S_f} f + \frac{2}{f}|\mathrm{d}f|^2_{\eta_{S_f}} \\ &= \frac{2}{f}(1 - \mathring{\Delta} \log f), \end{aligned} \tag{3.8}$$

where $\mathring{\Delta}$ is the Laplacian operator on the round sphere of radius 1. One can easily get the formula for $\mathrm{tr}\underline{\widetilde{\chi}}$ from the Gauss equation (2.8). Since the intrinsic geometry of S_f is conformal to the round

sphere with the conformal factor f^2, we have

$$K_{S_f} = f^{-2}(1 - \mathring{\Delta}\log f) = -\frac{1}{4}\mathrm{tr}\widetilde{\chi}\,\mathrm{tr}\underline{\widetilde{\chi}}, \tag{3.9}$$

which is consistent with the our formulas for $\mathrm{tr}\widetilde{\chi}$ and $\mathrm{tr}\underline{\widetilde{\chi}}$.

One can not find a closed trapped spacelike section in $\underline{C}_0$, since the Minkowski spacetime is complete. However as we already mentioned in the introduction that S. Klainerman, J. Luk and I. Rondianski could find a closed trapped surface in a vacuum spacetime which has sufficiently strong gravitation waves concentrated near some direction, it is interesting to see how the piece of the surface contained in the flat portion of the spacetime could be trapped.

It is well know that via the stereographic projection, one can conformally deform the round sphere minus a point to the flat plane. The conformal factor of the stereographic projection is e^{-w} where w is the Green's function for the Laplacian on the round sphere. In the spherical coordinates,

$$w = 2\log\sin\frac{\vartheta}{2}, \quad e^{-w} = \frac{1}{\sin^2\frac{\vartheta}{2}} = \frac{2}{1-\cos\vartheta}. \tag{3.10}$$

Since the Laplacian $\mathring{\Delta} = \frac{1}{\sin\vartheta}\partial_\vartheta(\sin\vartheta\partial_\vartheta) + \frac{1}{\sin^2\vartheta}\partial_\phi^2$, we have $\mathring{\Delta}\, w = -1$ outside the point $\vartheta = 0$. In the sense of distribution, w satisfies

$$\mathring{\Delta}\, w + 1 = 4\pi\delta, \tag{3.11}$$

where δ is the Dirac function at the point $\vartheta = 0$.

Considering the surface $\widetilde{S} = S_{e^{-w}}$, then its intrinsic metric $\eta|_{\widetilde{S}} = e^{-2w}(\mathrm{d}\vartheta^2 + \sin^2\vartheta\mathrm{d}\phi^2)$ has zero Gauss curvature. We have that

$$\mathrm{tr}\underline{\widetilde{\chi}} = -2e^w = \cos\vartheta - 1, \tag{3.12}$$

$$\mathrm{tr}\widetilde{\chi} = 0. \tag{3.13}$$

Hence $\widetilde{S}$ is marginally trapped. Easy to see that $\widetilde{S}$ is not compact, since it goes to infinity in the lightcone at the point $\vartheta = 0$.

Parameterized by $\{\vartheta, \phi\}$, we have the following map to represent $\widetilde{S}$ in the rectangular coordinates:

$$(\vartheta,\phi) \mapsto \left(\frac{2}{\cos\vartheta - 1}, \frac{2\sin\vartheta\cos\phi}{1-\cos\vartheta}, \frac{2\sin\vartheta\sin\phi}{1-\cos\vartheta}, \frac{2\cos\vartheta}{1-\cos\vartheta}\right). \tag{3.14}$$

Obviously, $\widetilde{S}$ lies in the null hyperplane $H_n = \{x_0 + x_3 = -2\}$. Actually $\widetilde{S} = \underline{C}_0 \cap H_n$, because by solving the following equation

$$\begin{cases} x_0 + x_3 = -2, \\ -x_0^2 + x_1^2 + x_2^2 + x_3^2 = 0, \end{cases} \tag{3.15}$$

we get exactly $\widetilde{S}$.

The intersection of $\underline{C}_0$ with the spacelike hyperplane $H_s = \{x_0 = -1\}$ is a spacelike surface $S' = \{(-1, \sin\vartheta\cos\phi, \sin\vartheta\sin\phi, \cos\phi)\}$. S' with its intrinsic metric is a round sphere of radius 1. The intersection of $\underline{C}_0$ with the timelike hyperplane $H_t = \{x_3 = -1\}$ is a spacelike surface $S'' = \{x_3 = -1, -x_0^2 + x_1^2 + x_2^2 = -1\} = \{\frac{1}{\cos\vartheta}, -\tan\vartheta\cos\phi, -\tan\vartheta\sin\phi, -1)\}$. S'' with its intrinsic metric is a hyperbolic surface of constant -1. S'' is in fact trapped, however it is not compact.

In fact the intersection of $\underline{C}_0$ with any space like hyperplane is intrinsically a round sphere, being of constant positive Gauss curvature, while the intersection with any timeline hyperplane is of constant negative Gauss curvature.

4 The short pulse hierarchy and the trapped surface formation theorem

In Christodoulou's monograph [1], he introduced the short pulse hierarchy and proved that the short pulse hierarchy is preserved in the evolution. The short pulse hierarchy is the key to the existence theorem as well as to the trapped surface formation theorem in his monograph. We present here the short pulse hierarchy and outline his argument of the trapped surface formation theorem.

Consider the characteristic initial data problem on an outgoing lightcone C. Let o be the vertex of the lightcone and T be a timelike vector at o. Then we can construct an affine parameter s on C such that the generator L of s satisfies that $g(L,T) = -1$ at o. We choose a constant $r_0 > 1$ and define the new affine parameter $\underline{u} = s - r_0$. We set the initial data before $\underline{u} = 0$ to be trivial. The initial data is nontrivial in the part $\{0 \leqslant \underline{u} \leqslant \delta\}$ for some small δ. Let ψ_0 be a smooth 2-dimensional symmetric trace-free matrix-valued function on $\mathbb{S}^2$, depending on $s \in [0,1]$, which extends smoothly by 0 to $s \leqslant 0$. We set

$$\psi(\underline{u},\theta) = \frac{\delta^{\frac{1}{2}}}{r_0}\psi_0(\frac{\underline{u}}{\delta},\theta), \quad (\underline{u},\theta) \in [0,\delta]\times\mathbb{S}^2. \tag{4.1}$$

Then the initial data in $0 \leqslant \underline{u} \leqslant \delta$ is given by the 2-covariant symmetric positive definite tensor desity m on $\mathbb{S}^2$ depending on $\underline{u}$

$$m = \exp\psi. \tag{4.2}$$

The incoming energy per unit solid angle in each direction in the interval $[0,\delta]$ is the integral

$$\frac{r_0^2}{8\pi}\int_0^\delta e\mathrm{d}\underline{u} \tag{4.3}$$

where

$$e = \frac{1}{2}|\widehat{\chi}|^2 = \frac{1}{8}|\partial_{\underline{u}}m|_m^2. \tag{4.4}$$

The factor $\delta^{\frac{1}{2}}$ in the ansatz (4.1) appears because of the desire to form trapped surfaces from the focusing of incoming waves.

By the theorem of domain of dependence, we know that the future M_0 of $\{\underline{u} \leqslant 0\}$ is flat. The future boundary of M_0 is the incoming lightcone $\underline{C}_{\underline{u}=0}$. Let Γ be the geodesic tangent to T and t be the affine parameter of Γ generated by T. We define a new function u along Γ by $u = \frac{t}{2} - r_0$ and u is extended as an optical function in the development which is constant along the future light cones with vertices on Γ. Set $u_0 = -r_0$. We are interested in the development of $\underline{C}_{\underline{u}=0,u_0\leqslant u\leqslant 0}$ and $C_{u=u_0,0\leqslant\underline{u}\leqslant\delta}$. In the development $\underline{u}$ is extended to be an optical function which is constant along the incoming null hypersurfaces corresponding to the level sets of $\underline{u}$ on C_{u_0}. Then the level sets of u and $\underline{u}$ give rise to a double null foliation in the development.

The full set of initial data is determined by integrating ordinary differential equations along the generators of C_{u_0}. The analysis of the equations along C_{u_0} gives, for the components of the spacetime curvature along C_{u_0}:

$$\begin{aligned}
\sup_{C_{u_0}} |\alpha| &\leqslant O_2(\delta^{-\frac{3}{2}}|u_0|^{-1}),\\
\sup_{C_{u_0}} |\beta| &\leqslant O_2(\delta^{-\frac{1}{2}}|u_0|^{-2}),\\
\sup_{C_{u_0}} |\rho|, \sup_{C_{u_0}} |\sigma| &\leqslant O_2(|u_0|^{-3}),\\
\sup_{C_{u_0}} |\underline{\beta}| &\leqslant O_2(\delta|u_0|^{-4}),\\
\sup_{C_{u_0}} |\underline{\alpha}| &\leqslant O_2(\delta^{\frac{3}{2}}|u_0|^{-5}),
\end{aligned} \tag{4.5}$$

The dependence on δ of the right hand sides of (4.5) displays the short pulse hierarchy.

Now we turn to Chirstodoulou's argument of the trapped surface formation theorem. Let $S_{u,\underline{u}}$ be the intersection of C_u with $\underline{C}_{\underline{u}}$ and H_{-1} be the spacelike hypersurface foliated by $\{S_{u,\underline{u}}\}_{u+\underline{u}=-1}$. The existence theorem guarantees that, for fixed r_0 and ψ_0, if we choose δ suitably small depending on r and ψ_0, we can solve the characteristic initial data problem and construct a regular spacetime at least up to H_{-1}. We denote by M_{-1} the spacetime bounded by C_{u_0}, $\underline{C}_0$, $\underline{C}_\delta$ and H_{-1}. In M_{-1}, the outmost spacelike section of the double null foliations is $S_{-1-\delta,\delta}$.

Let k be a constant greater than 1. If we assume that the incoming energy per unit solid angle in each direction in the advanced time interval $[0,\delta]$ is not less than $\frac{k}{8\pi}$, which is the uniform on $\mathbb{S}^2$ lower bound mentioned in the introduction, then for suitably small δ, $S_{-1-\delta,\delta}$ is actually trapped. For $\mathrm{tr}\underline{\chi}$, we have on $\underline{C}_\delta$

$$\left|\mathrm{tr}\underline{\chi} + \frac{2}{|u|}\right| \leqslant O(\delta|u|^{-2}). \tag{4.6}$$

For $\mathrm{tr}\chi$, we can estimate from the second variation formula along the generators of C_u and the incoming energy along C_u:

$$D\mathrm{tr}\chi = -\frac{1}{2}(\mathrm{tr}\chi)^2 - |\hat{\chi}|^2, \tag{4.7}$$

$$|u|^2|\widehat{\chi}|^2(u,\underline{u},\theta) \geqslant 2|u_0|^2 e(\underline{u},\theta) - O(\delta^{-\frac{1}{2}}). \tag{4.8}$$

We integrate the second variation formula (4.7) along the generators of C_u,

$$\mathrm{tr}\chi \leqslant \frac{2}{|u|} - \int_0^\delta |\widehat{\chi}|^2 \mathrm{d}\underline{u} = \frac{2}{|u|} - \frac{1}{|u|^2}\int_0^\delta |u|^2|\widehat{\chi}|^2 \mathrm{d}\underline{u}. \tag{4.9}$$

To estimate $\mathrm{tr}\chi$ through (4.9), we need to take the following fact into account that the integral in (4.9) is taken along the generators of C_u, not along the coordinate line with constant u and θ. Let $\{u, \underline{u}, \theta'\}$ be the double null coordinate system compatible with the generators of C_u and compatible with the generators of $\underline{C}_\delta$. Let p be any point of $\mathbb{S}^2$, we have

$$|u|^2|\widehat{\chi}|^2(u, \underline{u}, \theta' = p) \geqslant 2|u_0|^2 e(\underline{u}, \theta = p) - O(\delta^{-\frac{1}{2}}). \tag{4.10}$$

Then from (4.9) and (4.10), we get the following estimation of $\mathrm{tr}\chi$ on $\underline{C}_\delta$

$$\mathrm{tr}\chi(u, \delta, \theta') \leqslant \frac{2}{|u|} - \frac{2|u_0|^2}{|u|^2}\int_0^\delta e(\underline{u}, \theta = \theta')\mathrm{d}\underline{u} + O(\delta^{\frac{1}{2}}|u|^{-2}). \tag{4.11}$$

Hence if the incoming energy per unit solid angle in each direction in the advanced time interval $[0, \delta]$ is not less than $\frac{k}{8\pi}$, i.e.

$$\frac{|u_0|^2}{8\pi}\int_0^\delta e\mathrm{d}\underline{u} \geqslant \frac{k}{8\pi}, \tag{4.12}$$

we get

$$\mathrm{tr}\chi(u, \delta, \theta') \leqslant \frac{2}{|u|} - \frac{2k}{|u|^2} + O(\delta^{\frac{1}{2}}|u|^{-2}). \tag{4.13}$$

Then we conclude that if δ is suitably small,

$$\mathrm{tr}\chi(-1-\delta, \delta, \theta') \leqslant \frac{2 + 2\delta - 2k}{(1+\delta)^2} + O(\delta^{\frac{1}{2}}) < 0. \tag{4.14}$$

Hence $S_{-1-\delta,\delta}$ is trapped.

5 The transformation formula and the anisotropic criterion for formation of trapped surfaces

Klainerman, Luk and Rodianski can find a closed trapped surface $\widetilde{S}$ in $\underline{C}_\delta$ provided that the incoming energy per unit solid angle is large in some neighbourhood of some direction, even though in this case none of the spacelike section $S_{u,\underline{u}}$ of the double null foliations is trapped. The keys of their argument are a transformation formula for the null expansion $\mathrm{tr}\widetilde{\chi}$ of $\widetilde{S}$ and the Green's function of the Laplacian on the round sphere. We shall connect their argument with the material in section 3.

$\{u, \underline{u}, \theta\}$ is the double null coordinate system in M_{-1}. Any spacelike section $\widetilde{S}$ in $\underline{C}_\delta$ can be represented as the graph of a function f on $\mathbb{S}^2$, i.e. $\widetilde{S} = \{(u = -f(\theta), \delta, \theta)\}$. Then by the transformation formula for $\mathrm{tr}\widetilde{\chi}$ from [2],

$$\mathrm{tr}\widetilde{\chi} = \mathrm{tr}\chi - 2\Omega\Delta f - 4\Omega\eta\cdot\nabla f - 4\Omega^2\widehat{\underline{\chi}}(\nabla f, \nabla f) - \Omega^2\mathrm{tr}\underline{\chi}|\nabla f|^2 - 8\Omega^2\underline{\omega}|\nabla f|^2. \tag{5.1}$$

For a fixed function f on $\mathbb{S}^2$, as $\delta \to 0$, the transformation formula becomes

$$\mathrm{tr}\widetilde{\chi} = \mathrm{tr}\chi - \frac{2\overset{\circ}{\Delta} f}{f}. \tag{5.2}$$

Let $k(\theta)$ be the incoming energy per solid angle in the direction θ, i.e.

$$k(\theta) = \frac{r^2}{8\pi}\int_0^\delta e(\underline{u},\theta)\mathrm{d}\underline{u}. \tag{5.3}$$

Then we substitute the estimation (4.13) of $\mathrm{tr}\chi$ into the transformation formula (5.2) to get

$$\mathrm{tr}\widetilde{\chi}(\theta) \leqslant \frac{2}{f}(1-\overset{\circ}{\Delta}\log f) - \frac{2k(\theta)}{f^2} \tag{5.4}$$

as $\delta \to 0$. We shall use the knowledge about the sections of a lightcone in the Minkowski spacetime from section 3 to construct a function f such that the right hand side of (5.4) is negative, provided that k is large in some neighbourhood of some direction.

Let $\{\vartheta,\psi\}$ be the coordinate system on $\mathbb{S}^2$. We choose a positive smooth cutoff function g supported in $[0,2]$ such that $g \equiv 1$ on [0,1] and decreases to 0 on $[1,2]$. Let ε be an arbitrary small positive number. Denote $g_\varepsilon(\vartheta,\psi) = g(\frac{\vartheta}{\varepsilon})$. Then we construct f_ε by

$$\log f_\varepsilon = \begin{cases} -(1+\varepsilon)w, & \vartheta \in [2\varepsilon,\pi], \\ -(1-g_\varepsilon)(1+\varepsilon)w - g_\varepsilon(1+\varepsilon)w_\varepsilon, & \vartheta \in [0,2\varepsilon), \end{cases} \tag{5.5}$$

where $w(\vartheta,\psi) = 2\log\sin\frac{\vartheta}{2}$ is the Green's function of the Laplacian on the round sphere and $w_\varepsilon = 2\log\sin\frac{\varepsilon}{2}$.

For $\vartheta \in [2\varepsilon,\pi]$, $\overset{\circ}{\Delta}\log f_\varepsilon = -(1+\varepsilon)\overset{\circ}{\Delta} w = 1+\varepsilon$ and $f_\varepsilon(\pi) = 1$. Let k_ε in the following be:

$$k_\varepsilon = \sup_{\vartheta\in[0,2\varepsilon]} f_\varepsilon(1-\overset{\circ}{\Delta}\log f_\varepsilon) \tag{5.6}$$

We choose initial data such that the incoming energy per unit solid angle $k(\vartheta,\psi)$ in $\{\vartheta\in[0,2\varepsilon]\}$ is larger than k_ε. In the following we prove that for δ suitable small, the surface $S_{f_\varepsilon} = \{(-f_\varepsilon(\theta),\delta,\theta)\}$ is trapped.

For $\vartheta \in [0,2\varepsilon]$,

$$\begin{aligned} \mathrm{tr}\widetilde{\chi} &\leqslant \frac{2}{f_\varepsilon}(1-\overset{\circ}{\Delta}\log f_\varepsilon) - \frac{2k(\vartheta,\psi)}{f_\varepsilon^2} \\ &\leqslant \frac{2k_\varepsilon}{f_\varepsilon^2} - \frac{2k(\vartheta,\psi)}{f_\varepsilon^2} \\ &< 0. \end{aligned} \tag{5.7}$$

For $\vartheta \in [2\varepsilon, \pi]$,

$$\begin{aligned} \mathrm{tr}\widetilde{\chi} &\leqslant \frac{2}{f_\varepsilon}(1 - \overset{\circ}{\Delta} \log f_\varepsilon) - \frac{2k(\vartheta,\psi)}{f_\varepsilon^2} \\ &\leqslant -\frac{2\varepsilon}{f_\varepsilon} \\ &< 0. \end{aligned} \tag{5.8}$$

Hence $\mathrm{tr}\widetilde{\chi}$ is negative on S_{f_ε} for δ suitable small.

For the null expansion $\mathrm{tr}\underline{\widetilde{\chi}}$ on S_{f_ε}, since $\mathrm{tr}\underline{\widetilde{\chi}} = \mathrm{tr}\underline{\chi}$ and on $\underline{C}_\delta$,

$$\left| \mathrm{tr}\underline{\chi} + \frac{2}{|u|} \right| \leqslant O(\delta |u|^{-2}), \tag{5.9}$$

we get as $\delta \to 0$,

$$\mathrm{tr}\underline{\widetilde{\chi}} = -\frac{2}{f_\varepsilon} < 0. \tag{5.10}$$

Hence $\mathrm{tr}\underline{\widetilde{\chi}}$ is negative on S_{f_ε} for δ suitable small. Hence S_{f_ε} is trapped.

Now we can estimate the size of the trapped surface S_{f_ε}. In the direction $\vartheta = \pi$, S_{f_ε} is near to the section $S_{-1-\delta,\delta}$ of the double null foliation, since $f_\vartheta = 1$ at $\vartheta = \pi$. In the direction $\vartheta = 0$, S_{f_ε} is far away from the section $S_{-1-\delta,\delta}$ of the double null foliation, since $f_\varepsilon = \left(\frac{2}{1-\cos\varepsilon}\right)^{1+\varepsilon}$ at $\vartheta = 0$ and $\left(\frac{2}{1-\cos\varepsilon}\right)^{1+\varepsilon}$ is much greater than 1 for small ε. As $\varepsilon \to 0$, $\varepsilon^2 f_\varepsilon$ tends to a finite limit, i.e. $f_\varepsilon \sim \varepsilon^{-2}$.

We can also obtain a upper bound of k_ε. In the coordinate system $\{\vartheta, \psi\}$, $\overset{\circ}{\Delta} = \partial_\vartheta^2 + \cot\vartheta\partial_\vartheta + \sin^{-2}\vartheta\partial_\psi^2$. Then for $\vartheta \in [2\varepsilon, \varepsilon]$,

$$w = O(\log\varepsilon), \quad \partial_\vartheta g_\varepsilon = O(\varepsilon^{-1}), \quad \partial_\vartheta w = O(\varepsilon^{-1}), \quad \overset{\circ}{\Delta} g_\varepsilon = O(\varepsilon^{-2}), \quad \overset{\circ}{\Delta} w = O(\varepsilon^{-2}). \tag{5.11}$$

Hence $\overset{\circ}{\Delta} \log f_\varepsilon = O(\varepsilon^{-2}\log\varepsilon)$ for $\vartheta \in [\varepsilon, 2\varepsilon]$. For $\vartheta \in [0, \varepsilon]$, $\overset{\circ}{\Delta} \log f_\varepsilon = 0$ since $\log f_\varepsilon$ is constant. So for $\vartheta \in [0, 2\varepsilon]$, $\overset{\circ}{\Delta} \log f_\varepsilon = O(\varepsilon^{-2}\log\varepsilon)$. Then we conclude that $k_\varepsilon = O(\varepsilon^{-4}\log\varepsilon)$.

References

[1] Christodoulou, D., *The Formation of Black Holes in General Relativity*. EMS Monographs in Mathematics, EMS Publishing House, 2007.

[2] Klainerman, S., Luk, J., Rodnianski, I., A fully anisotropic mechanism for formation of trapped surfaces in vacuum, *Invent. Math.* **198**, 1–26

[3] Penrose, R., Gravitational collapse and space-time singularitie, *Phys. Rev. Lett.* **14** (1965), 57–59.